

Clinical Policy: Lumacaftor/Ivacaftor (Orkambi)

Reference Number: CP.PHAR.213

Effective Date: 05.01.16

Last Review Date: 02.20

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Lumacaftor/ivacaftor (Orkambi[®]) is a combination drug for cystic fibrosis (CF). Lumacaftor improves the conformational stability of F508del-cystic fibrosis transmembrane conductance regulator (CFTR), while ivacaftor is a CFTR potentiator.

FDA Approved Indication(s)

Orkambi is indicated for the treatment of CF in patients age 2 years and older who are homozygous for the F508del mutation in the CFTR gene.

If the patient's genotype is unknown, an FDA-cleared CF mutation test should be used to detect the presence of the F508del mutation on both alleles of the CFTR gene.

Limitation(s) of use: The efficacy and safety of Orkambi have not been established in patients with CF other than those homozygous for the F508del mutation.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Orkambi is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Cystic Fibrosis (must meet all):

1. Diagnosis of CF confirmed by all of the following (a, b, and c):
 - a. Clinical symptoms consistent with CF in at least one organ system, or positive newborn screen or genetic testing for siblings of patients with CF;
 - b. Evidence of CFTR dysfunction confirmed by one of the following (i or ii) (*see Appendix D*):
 - i. Elevated sweat chloride ≥ 60 mmol/L;
 - ii. Genetic testing confirming the presence of two disease-causing mutations in CFTR gene, one from each parental allele;
 - c. Member is homozygous for the *F508del* mutation in the CFTR gene;
2. Age ≥ 2 years;
3. Prescribed by or in consultation with a pulmonologist;

4. Chart notes indicate that pulmonary function tests, performed within the last 90 days, show a percent predicted forced expiratory volume in 1 second (ppFEV1) that is between 40-90%;
5. Orkambi is not prescribed concurrently with other CFTR modulators (e.g., Kalydeco[®], Symdeko[®], Trikafta[™]);
6. Dose does not exceed one of the following (a, b, c, or d):
 - a. Age 2 to 5 years weighing < 14 kg: lumacaftor 200 mg/ivacaftor 250 mg (2 packets) per day;
 - b. Age 2 to 5 years weighing ≥ 14 kg: lumacaftor 300 mg/ivacaftor 376 mg (2 packets) per day;
 - c. Age 6 to 11 years: lumacaftor 400 mg/ivacaftor 500 mg (4 tablets) per day;
 - d. Age ≥ 12 years: lumacaftor 800 mg/ivacaftor 500 mg (4 tablets) per day.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Cystic Fibrosis (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by a stabilization in ppFEV1 if baseline was ≥ 70% or increase in ppFEV1 if baseline was < 70%;
3. Orkambi is not prescribed concurrently with other CFTR modulators (e.g., Kalydeco, Symdeko, Trikafta);
4. If request is for a dose increase, new dose does not exceed one of the following (a, b, c, or d):
 - a. Age 2 to 5 years weighing < 14 kg: lumacaftor 200 mg/ivacaftor 250 mg (2 packets) per day;
 - b. Age 2 to 5 years weighing ≥ 14 kg: lumacaftor 300 mg/ivacaftor 376 mg (2 packets) per day;
 - c. Age 6 to 11 years: lumacaftor 400 mg/ivacaftor 500 mg (4 tablets) per day;
 - d. Age ≥ 12 years: lumacaftor 800 mg/ivacaftor 500 mg (4 tablets) per day.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CF: cystic fibrosis

CFTR: cystic fibrosis transmembrane conductance regulator

FDA: Food and Drug Administration

ppFEV1: percent predicted forced expiratory volume in 1 second

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: General Information

- Regarding the diagnostic criteria for CF of “genetic testing confirming the presence of two disease-causing mutations in CFTR gene,” this is to ensure that whether heterozygous or homozygous, there are two disease-causing mutations in the CFTR gene, one from each parental allele.
- Most children can do spirometry by age 6, though some preschoolers are able to perform the test at a younger age. Some young children aren’t able to take a deep enough breath and blow out hard and long enough for spirometry. Forced oscillometry is another way to test lung function in young children. This test measures how easily air flows in the lungs (resistance and compliance) with the use of a machine.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
CF	Adults and pediatric patients age 12 years and older: two tablets (each containing lumacaftor 200 mg/ivacaftor 125 mg) PO Q12H	Adults and pediatric patients age 12 years and older: lumacaftor 800 mg/ivacaftor 500 mg per day
	Pediatric patients age 6 through 11 years: two tablets (each containing lumacaftor 100 mg/ivacaftor 125 mg) PO Q12H	Pediatric patients age 6 through 11 years: lumacaftor 400 mg/ivacaftor 500 mg per day
	Pediatric patients age 2 through 5 years and weighing < 14 kg: one packet of granules (each containing lumacaftor 100 mg/ivacaftor 125 mg) PO Q12H	Pediatric patients age 2 through 5:

Indication	Dosing Regimen	Maximum Dose
	Pediatric patients age 2 through 5 years and weighing ≥ 14 kg: one packet of granules (each containing lumacaftor 150 mg/ivacaftor 188 mg) PO Q12H	<14 kg - lumacaftor 200 mg/ivacaftor 250 mg per day ≥ 14 kg - lumacaftor 300 mg/ivacaftor 376 mg per day

VI. Product Availability

- Tablets: lumacaftor 100 mg and ivacaftor 125 mg, lumacaftor 200 mg and ivacaftor 125 mg
- Oral granules: lumacaftor 100 mg and ivacaftor 125 mg, lumacaftor 150 mg and ivacaftor 188 mg

VII. References

1. Orkambi Prescribing Information. Boston, MA: Vertex Pharmaceuticals, Inc.; July 2019. Available at <http://www.orkambi.com>. Accessed October 28, 2019.
2. Mogayzel PJ, Naureckas ET, Robinson KA, et al. Cystic fibrosis pulmonary guidelines: Chronic medications for maintenance of lung health. Am J Respir Crit Care Med. 2013; 187(7): 680-689.
3. Farrell PM, White TB, Ren CL et al. Diagnosis of cystic fibrosis: Consensus guidelines from the Cystic Fibrosis Foundation. J Pediatr. 2017; 181S: S4-15.
4. Ren CL, Morgan RL, Oermann C, et al. Cystic Fibrosis Foundation pulmonary guidelines: Use of cystic fibrosis transmembrane conductance regulator modulator therapy in patients with cystic fibrosis. Ann Am Thorac Soc. 2018; 15(3): 271-280.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy split from CP.PHAR.54 CF Treatments. Evidence of a “significant improvement in FEV1” to continue approval is replaced with “Demonstrated positive response (improvement, maintenance, decreased rate of progression/decline) to Orkambi therapy in one or more of the following areas: pulmonary function, quality of life, pulmonary exacerbations”. Not having increased LFTs is removed as a discontinuation reason. Continuation approval period is extended from 6 to 12 months.	04.16	05.16
Age lowered to 6 years per PI – corresponding maximum dose added. Efficacy statement edited to indicate general positive response to therapy.	05.17	05.17
1Q18 annual review: - Policies combined for Centene Medicaid and Commercial lines of business. - No significant changes from previous corporate approved policy - Commercial: Added age requirement per FDA labeling. Modified max dose criteria to be age-specific	10.26.17	02.18

Reviews, Revisions, and Approvals	Date	P&T Approval Date
- References reviewed and updated.		
No significant changes: updated age limit with corresponding dosing for pediatric patients down to 2 years of age per updated prescribing information.	09.26.18	
1Q 2019 annual review: no significant changes; references reviewed and updated.	10.25.18	02.19
Added HIM line of business per SDC and prior approved clinical guidance.	04.01.19	
1Q 2020 annual review: added the following criteria to initial approval: comprehensive diagnostic criteria (e.g., clinical symptoms in at least one organ, positive newborn screen, siblings genetic testing, and evidence of CFTR dysfunction) to confirm diagnosis of CF, prescriber requirement of pulmonologist, chart notes indicate that pulmonary function tests (ppFEV1 between 40-90%), not prescribed concurrently with other CFTR modulators; added the following to continued therapy criteria: positive response as evidenced by stabilization in ppFEV1 in lieu of an increase is acceptable if baseline was $\geq 70\%$, not prescribed concurrently with other CFTR modulators; added Appendix D; changed approval durations of commercial from length of benefit to 6 months initial and 12 months continued; references reviewed and updated.	12.17.19	02.20

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.