

Clinical Policy: Neratinib (Nerlynx)

Reference Number: CP.PHAR.365

Effective Date: 12.01.17 Last Review Date: 05.20

Line of Business: Commercial, HIM, Medicaid Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Neratinib (Nerlynx[®]) is a kinase inhibitor that irreversibly binds to epidermal growth factor receptor, human epidermal growth factor receptor 2 (HER2), and HER4.

FDA Approved Indication(s)

Nerlynx is indicated:

- As a single agent, for the extended adjuvant treatment of adult patients with early stage HER2-overexpressed/amplified breast cancer, to follow adjuvant trastuzumab-based therapy.
- In combination with capecitabine, for the treatment of adult patients with advanced or metastatic HER2-positive breast cancer who have received two or more prior anti-HER2 based regimens in the metastatic setting.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Nerlynx is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Breast Cancer (must meet all):
 - 1. Diagnosis of breast cancer;
 - 2. Prescribed by or in consultation with an oncologist;
 - 3. Age \geq 18 years;
 - 4. Disease is HER2-positive;
 - 5. Member meets one of the following (a or b):
 - a. Both (i and ii):
 - i. Documentation of previous adjuvant treatment with trastuzumab;
 - ii. Disease is early stage (stage 1-3) or hormone-receptor positive;
 - b. Prescribed in combination with capecitabine for recurrent, advanced, or metastatic disease, and member has received two or more prior anti-HER2 based regimens used in the metastatic setting;
 - 6. Request meets one of the following (a or b):*
 - a. Dose does not exceed 240 mg (6 tablets) per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

^{*}Prescribed regimen must be FDA-approved or recommended by NCCN

CLINICAL POLICY Neratinib



Approval duration: Medicaid/HIM – 6 months Commercial – Length of Benefit

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Breast Cancer (member meets all):

- 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Nerlynx for breast cancer and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 240 mg (6 tablets) per day;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 12 months

Commercial – Length of Benefit

B. Other diagnoses/indications (member meets 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration

HER: human epidermal growth factor receptor NCCN: National Comprehensive Cancer Network





Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business

and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/
Herceptin [®]	Administer according to one of the following doses	Maximum Dose 8 mg/kg
(trastuzumab)	and schedules for a total of 52 weeks:	8 8
Ogivri TM	Herceptin, Ogivri, Herzuma, Ontruzant, Trazimera,	
(trastuzumab-	Kanjinti:	
dkst)	During and following paclitaxel, docetaxel, or	
Ontruzant [®]	docetaxel/carboplatin:	
(Trastuzumab-	• Initial dose of 4 mg/kg as an IV infusion over 90	
dttb)	minutes then at 2 mg/kg as an IV infusion over	
Herzuma®	30 minutes weekly during chemotherapy for the	
(Trastuzumab-	first 12 weeks (paclitaxel or docetaxel) or 18	
pkrb)	weeks (docetaxel/carboplatin).	
Trazimera TM	One week following the last weekly dose of the	
(Trastuzumab-	trastuzumab product, administer trastuzumab	
qyyp)	product at 6 mg/kg as an IV infusion over 30 to	
Kanjinti™	90 minutes every 3 weeks.	
(Trastuzumab-	Herceptin, Ogivri, Ontruzant, Trazimera, Kanjinti:	
anns)	As a single agent within 3 weeks following	
,	completion of multi-modality, anthracycline based	
	chemotherapy regimens:	
	• Initial dose: 8 mg/kg as an IV infusion over 90	
	minutes.	
	• Subsequent doses: 6 mg/kg as an IV infusion	
	over 30 to 90 minutes every 3 weeks	
Herceptin	Herceptin Hylecta (subcutaneous product):	600 mg/10,000
Hylecta TM	As part of a treatment regimen consisting of	units every 3
(Trastuzumab-	doxorubicin, cyclophosphamide, and either paclitaxel	weeks
hyaluronidase-	or docetaxel; as part of a treatment regimen with	
oysk)	docetaxel and carboplatin; as a single agent following	
<i>3</i>	multi-modality anthracycline based therapy: 600 mg	
	trastuzumab and 10,000 units hyaluronidase	
	administered subcutaneously over approximately 2-5	
	minutes once every 3 weeks	
Perjeta®	Initial 840 mg IV followed by a maintenance dose of	Maintenance:
(pertuzumab)	420 mg IV every 3 weeks in combination with	420 mg every 3
, <u> </u>	trastuzumab and either docetaxel or paclitaxel	weeks
Kadcyla® (ado-	3.6 mg/kg IV every 3 weeks	3.6 mg/kg every
trastuzumab		3 weeks
emtansine)		
Enhertu® (fam-	5.4 mg/kg once every 3 weeks	5.4 mg/kg every
trastuzumab		3 weeks



Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
deruxtecan- nxki)		

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings None reported

Appendix D: General Information

- Per the Nerlynx prescribing information, antidiarrheal prophylaxis is recommended during the first 2 cycles (56 days) of Nerlynx treatment and should be initiated with the first dose of Nerlynx in order to address the risk of treatment discontinuation due to diarrhea, as was seen in the pivotal ExteNET trial.
- Nerlynx is FDA-approved for a one year total duration of therapy as it was only administered for one year in the pivotal ExteNET trial; however, the NCCN does not recommend any specific length of treatment.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Breast cancer extended	240 mg PO QD	240 mg/day
adjuvant treatment		
Breast cancer advanced,	240 mg PO QD on days 1-21 plus	240 mg/day
recurrent, or metastatic	capecitabine 750 mg/m ² PO BID on	
disease	days 1-14 of a 21-day cycle	

VI. Product Availability

Tablet: 40 mg

VII. References

- 1. Nerlynx Prescribing Information. Los Angeles, CA: Puma Biotechnology, Inc.; February 2020. Available at: www.nerlynx.com. Accessed April 7, 2020.
- 2. National Comprehensive Cancer Network. Breast Cancer Version 3.2020. Available at: https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed April 7, 2020.
- 3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug compendium. Accessed April 7, 2020.
- 4. National Comprehensive Cancer Network. Central Nervous System Cancers Version 1.2020. Available at: https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf. Accessed April 7, 2020.

Reviews, Revisions, and Approvals	Date	P&T
		Approval Date
Policy created	09.05.17	11.17
Combined Commercial and Medicaid policies.	02.12.18	



Reviews, Revisions, and Approvals	Date	P&T
		Approval
		Date
4Q 2018 annual review: added NCCN off-label uses; added	07.06.18	11.18
specialist involvement in care; removed restriction for only 1 year		
of total therapy as NCCN does not recommend a specific duration		
of use; references reviewed and updated.		
4Q 2019 annual review: no significant changes; removed off-label	08.12.19	11.19
capecitabine combination use from criteria (NCCN category 2B);		
references reviewed and updated.		
RT2: added new indication for use in combination with	04.07.20	05.20
capecitabine for advanced, recurrent, or metastatic breast cancer;		
added HIM line of business; added Commercial length of benefit		
authorization for initial and continuation of therapy.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise

CLINICAL POLICY Neratinib



professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation. The composition of the contained herein composition.