

Clinical Policy: Ibrutinib (Imbruvica)

Reference Number: CP.PHAR.126

Effective Date: 10.01.15

Last Review Date: 02.22

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Ibrutinib (Imbruvica[®]) is a Bruton tyrosine kinase (BTK) inhibitor.

FDA Approved Indication(s)

Imbruvica is indicated for the treatment of:

- Adult patients with mantle cell lymphoma (MCL) who have received at least one prior therapy*
- Adult patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL)
- Adult patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) with 17p deletion
- Adult patients with Waldenström's macroglobulinemia (WM)
- Adult patients with marginal zone lymphoma (MZL) who require systemic therapy and have received at least one prior anti-CD20-based therapy*
- Adult patients with chronic graft-versus-host disease (cGVHD) after failure of one or more lines of systemic therapy

*Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Imbruvica is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Mantle Cell Lymphoma (*B-cell lymphoma subtype*) (must meet all):**

1. Diagnosis of MCL (a B-cell lymphoma subtype);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Member meets one of the following* (a or b):

- a. Prescribed in combination with rituximab as pretreatment in order to limit the number of cycles of aggressive induction therapy with RHyperCVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone);
 - b. Received ≥ 1 prior line of systemic therapy (*see Appendix B*);
**Prior authorization may be required*
6. Request meets one of the following (a, b, or c):*
- a. For dose ≤ 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - b. For dose ≥ 420 mg (not to exceed 560 mg) per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

B. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (must meet all):

1. Diagnosis of CLL or SLL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age ≥ 18 years;
4. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Prescribed as a single agent; or in combination with one of the following* (a, b, or c):
 - a. Rituxan[®] (rituximab);
 - b. Gazyva[®] (obinutuzumab);
 - c. Bendamustine and Rituxan;**Prior authorization may be required*
6. Request meets one of the following (a, b, or c):*
 - a. For dose ≤ 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - b. For 420 mg dose per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

C. Waldenström's Macroglobulinemia (must meet all):

1. Diagnosis of WM or lymphoplasmacytic lymphoma;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age ≥ 18 years;
4. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Prescribed as a single agent or in combination with Rituxan*;

**Prior authorization may be required*

6. Request meets one of the following (a, b, or c):*
 - a. For dose \leq 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - b. For 420 mg dose per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

D. Marginal Zone Lymphoma (*B-cell lymphoma subtype*) (must meet all):

1. Diagnosis of one of the following MZL subtypes (a, b, c, or d):
 - a. Gastric MALT lymphoma;
 - b. Nongastric MALT lymphoma (noncutaneous);
 - c. Nodal MZL;
 - d. Splenic MZL;
 2. Prescribed by or in consultation with an oncologist or hematologist;
 3. Age \geq 18 years;
 4. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
 5. Received \geq 1 line of systemic therapy* (*see Appendix B*);
- *Prior authorization may be required*
6. Request meets one of the following (a, b, or c):*
 - a. For dose \leq 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - b. For dose \geq 420 mg (not to exceed 560 mg) per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

E. Chronic Graft-Versus-Host Disease (must meet all):

1. Diagnosis of cGVHD;
2. Prescribed by or in consultation with an oncologist, hematologist, or bone marrow transplant specialist;
3. Age \geq 18 years;
4. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Member has a history of bone marrow/stem cell transplant;
6. Member meets one of the following (a and b):

- a. Failure of a systemic corticosteroid (*see Appendix B*) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
- b. Failure of a systemic immunosuppressant (*see Appendix B*) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required*

7. Imbruvica is not prescribed concurrently with Jakafi[®] or Rezurock[™];
8. Request meets one of the following (a, b, or c):*
 - a. For dose \leq 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - b. For 420 mg dose per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

F. NCCN Compendium Indications (off-label) (must meet all):

1. Diagnosis of one of the following (a, b, or c):
 - a. Primary CNS lymphoma;
 - b. Hairy cell leukemia (HCL);
 - c. B-cell lymphoma subtype (i, ii, iii, iv, v, or vi):
 - i. AIDS-related non-germinal center DLBCL;
 - ii. High-grade B-cell lymphoma;
 - iii. Follicular lymphoma (grade 1-2) (FL);
 - iv. Post-transplant lymphoproliferative disorder (PTLD);
 - v. DLBCL;
 - vi. Histologic transformation of MZL to DLBCL;
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age \geq 18 years;
4. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
5. Member meets one of the following (a or b):
 - a. For primary CNS lymphoma, request is for use as either induction therapy or for relapsed or refractory disease;
 - b. For B-cell lymphoma, received \geq 1 prior line of systemic therapy (*see Appendix B*), unless contraindicated or clinically significant adverse effects are experienced to all;
 - c. For HCL, received \geq 2 prior lines of systemic therapies (*see Appendix B*), unless contraindicated or clinically significant adverse effects are experienced to all;
6. Request meets one of the following (a, b, or c):*
 - a. For dose \leq 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;

- b. For dose ≥ 420 mg (not to exceed 560 mg) per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
- c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 6 months

Commercial – Length of Benefit

G. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Imbruvica for a covered oncology-related indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. For Imbruvica requests, member must use ibrutinib, if available, unless contraindicated or clinically significant adverse effects are experienced;
4. For cGVHD, Imbruvica is not prescribed concurrently with Jakafi[®] or Rezurock[™];
5. If request is for a dose increase, request meets one of the following (a, b, or c):*
 - a. MCL and MZL, new dose does not exceed 560 mg per day and one of the following (i or ii):
 - i. For dose ≤ 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - ii. For dose ≥ 420 mg (not to exceed 560 mg) per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - b. CLL/SLL, WM, and cGVHD, new dose does not exceed 420 mg and one of the following (i or ii):
 - i. For dose ≤ 420 mg per day, request is for capsules and prescribed quantity does not exceed 3 capsules per day;
 - ii. For 420 mg dose per day, request is for tablets and prescribed quantity does not exceed 1 tablet per day;
 - c. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**For oncology indications, prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration:

Medicaid/HIM – 12 months

Commercial – Length of Benefit

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

- Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

BTK: Bruton’s tyrosine kinase	MCL: mantle cell lymphoma
cGVHD: chronic graft-versus-host disease	MZL: marginal zone lymphoma
CLL: chronic lymphocytic leukemia	NCCN: National Comprehensive Cancer Network
DLBCL: diffuse large B-cell lymphoma	PTLD: post-transplant lymphoproliferative disorders
FDA: Food and Drug Administration	SLL: small lymphocytic lymphoma
FL: follicular lymphoma	WM: Waldenström’s macroglobulinemia
HCL: hairy cell leukemia	
MALT: mucosa-associated lymphoid tissue	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<i>Examples of systemic therapies for B-cell lymphomas</i>		
Bendeka [®] , Treanda [®] (bendamustine) ± Rituxan (rituximab) or Gazyva [®] (obinutuzumab)	Varies	Varies
CHOP + Gazyva (obinutuzumab)		
EPOCH [etoposide, prednisone, vincristine (Vincasar PFS [®]), cyclophosphamide, doxorubicin (Adriamycin [®])] + Rituxan (rituximab)		
NORDIC [dose-intensified induction immunochemotherapy with Rituxan (rituximab) + cyclophosphamide, vincristine (Vincasar PFS), doxorubicin, prednisone] alternating with Rituxan (rituximab) and high-dose cytarabine		
RCEOP [Rituxan (rituximab), cyclophosphamide, etoposide, vincristine (Vincasar PFS), prednisone]		
RCEPP [Rituxan (rituximab), cyclophosphamide, etoposide, prednisone, procarbazine]		

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
RCHOP [cyclophosphamide, doxorubicin (Adriamycin [®]), vincristine (Vincasar PFS), prednisone]/RDHAP		
RCVP [Rituxan (rituximab), cyclophosphamide, doxorubicin (Adriamycin [®]), vincristine (Vincasar PFS)]		
RDHAP [Rituxan (rituximab), dexamethasone, cytarabine, cisplatin]		
RDHAX [Rituxan (rituximab), dexamethasone, cytarabine, oxaliplatin]		
Revlimid [®] (lenalidomide) + Rituxan (rituximab)		
Rituxan (rituximab)		
VR-CAP [bortezomib (Velcade [®]), Rituxan (rituximab), cyclophosphamide, doxorubicin (Adriamycin [®]), and prednisone]		
Examples of systemic corticosteroids and immunosuppressants for cGVHD		
Systemic corticosteroids (e.g., methylprednisolone, prednisone)	Varies	Varies
mycophenolate mofetil (Cellcept [®])		
cyclosporine (Gengraf [®] , Neoral [®] , Sandimmune [®])		
tacrolimus (Prograf [®])		
sirolimus (Rapamune [®])		
imatinib (Gleevec [®])		
Jakafi [®] (ruxolitinib)		
Rezurock [™] (belumosudil)		
Examples of systemic therapies for primary CNS lymphoma		
High-dose methotrexate-based regimen [methotrexate (Rheumatrex [®]) + Rituxan (rituximab) and other agents (e.g., temozolomide, vincristine (Vincasar PFS), procarbazine, cytarabine)]	Varies	Varies
Examples of systemic therapies for HCL		
Intron [®] A (interferon alfa-2b)	Varies	Varies
cladribine		
Nipent [™] (pentostatin)		

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
MCL, MZL	560 mg PO QD	560 mg/day (3 capsules or 1 tablet per day)
CLL/SLL, WM, cGVHD	420 mg PO QD	420 mg/day (3 capsules or 1 tablet per day)

VI. Product Availability

- Capsules: 70 mg, 140 mg
- Tablets: 140 mg, 280 mg, 420 mg, 560 mg

VII. References

1. Imbruvica Prescribing Information. Sunnyvale, CA: Pharmacyclics LLC; December 2020. Available at: <https://www.imbruvica.com/>. Accessed November 13, 2021.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed November 9, 2020.
3. National Comprehensive Cancer Network. Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Version 1.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/cll.pdf. Accessed November 13, 2021.
4. National Comprehensive Cancer Network. Hairy Cell Leukemia Version 1.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/hairy_cell.pdf. Accessed November 13, 2021.
5. National Comprehensive Cancer Network. Central Nervous System Cancers Version 2.2021. Available at: https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf. Accessed November 13, 2021.
6. National Comprehensive Cancer Network. Waldenström’s Macroglobulinemia/Lymphoplasmacytic Lymphoma Version 1.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/waldenstroms.pdf. Accessed November 13, 2021.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Converted to new template. Added new FDA approved indication: cGVHD. Increased continued approval duration from 6 to 12 months. Created criteria for hairy cell leukemia per NCCN guidelines/compendium. Added Appendix B: General Information.	08.09.17	11.17
3Q 2018 annual review: Policies combined for commercial, HIM, and Medicaid lines of business; For all lines of business: off-label NCCN compendium-supported uses were added, tablet formulations were added, age requirement was added for FDA-labeled indications, specialist requirement was added for all indications; For commercial: added off-label use of ibrutinib pretreatment for MCL per NCCN guidelines; For Medicaid, removed age requirement for pretreatment	05.15.18	08.18

Reviews, Revisions, and Approvals	Date	P&T Approval Date
use of ibrutinib for MCL per NCCN guidelines; references reviewed and updated.		
Per SDC, added preferencing for capsule formulation.	10.05.18	
1Q 2019 annual review: for CLL/SLL, added requirement for single agent use per updated NCCN guidelines since combo use is category 2B; for FL, revised requirement of trial and failure to one prior therapy instead of two per updated NCCN guidelines; for CNS lymphoma, added hematologist prescriber option; consolidated criteria for NCCN compendium off-label uses; references reviewed and updated.	11.06.18	02.19
1Q 2020 annual review: no significant changes; references reviewed and updated.	11.26.19	02.20
RT4: modified CLL/SLL and WM criteria to allow combination use per updated FDA labeling (indication language remains unchanged). Revised maximum quantity by dose to maximize dose form cost effectiveness per data analytics recommendation; removed requirement for medical justification why capsules cannot be used.	04.28.20	
1Q 2021 annual review: oral oncology generic redirection language added; for MCL, NCCN directed language inserted to clarify combination therapy with rituximab; for CLL/SCC, histologic transformation combination therapy added per NCCN; for MZL, subtypes delineated for clarity, therapy trials broadened beyond rituximab per NCCN; for cGVHD, trial requirement edited to require a systemic corticosteroid and an immunosuppressant agent per NCCN and the Imbruvica pivotal trial; Appendix B reorganized by B-cell lymphomas vs. other indications; references to HIM.PHAR.21 to revised to HIM.PA.154; references reviewed and updated.	11.09.20	02.21
Added language for Imbruvica, Rezerox and Jafaki not to be used concurrently since all are used for cGVHD. Updated Appendix B alternatives for cGVHD.	08.24.21	11.21
1Q 2022 annual review: removed indication for CLL/SLL histologic (Richter's) transformation per NCCN as it is now category 2B; added indication of lymphoplasmacytic lymphoma to WM criteria per NCCN; updated primary CNS lymphoma criterion that ibrutinib may be used as either induction therapy or for relapsed or refractory disease per NCCN; clarified oral oncology generic redirection language to “must use”; references reviewed and updated.	11.13.21	02.22

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional

organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2015 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.