

Clinical Policy: Deferasirox (Exjade, Jadenu)

Reference Number: CP.PHAR.145

Effective Date: 11.01.15

Last Review Date: 08.21

Line of Business: Commercial, HIM, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Deferasirox (Exjade[®], Jadenu[®]) is an iron chelator.

FDA Approved Indication(s)

Exjade and Jadenu are indicated for the treatment of:

- Chronic iron overload due to blood transfusions (transfusional hemosiderosis) in patients 2 years of age and older.
- Chronic iron overload in patients 10 years of age and older with non-transfusion-dependent thalassemia (NTDT) syndromes and with a liver iron concentration (LIC) of at least 5 milligrams of iron per gram of liver dry weight (mg Fe/g dw) and a serum ferritin greater than 300 mcg/L.

Limitation(s) of use:

The safety and efficacy of Exjade/Jadenu when administered with other iron chelation therapy have not been established.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Exjade and Jadenu are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Chronic Iron Overload due to Blood Transfusions (must meet all):

1. Diagnosis of chronic iron overload due to blood transfusions;
2. Age \geq 2 years;
3. Member must use generic deferasirox, unless contraindicated or clinically significant adverse effects are experienced;
4. Transfusion history of \geq 100 mL/kg of packed red blood cells (e.g., \geq 20 units of packed red blood cells for a 40 kg person) and a serum ferritin level $>$ 1,000 mcg/L;
5. At the time of the request, member has none of the following contraindications:
 - a. Glomerular filtration rate (GFR) $<$ 40 mL/min/1.73 m²;
 - b. Platelet count $<$ 50 x 10⁹/L;
 - c. Severe hepatic impairment (Child-Pugh C);
6. Therapy does not include concurrent use of other iron chelators;
7. Dose does not exceed the following (a or b):

- a. Exjade: 40 mg/kg per day (*see Appendix D for dose rounding guidelines*);
- b. Jadenu: 28 mg/kg per day (*see Appendix D for dose rounding guidelines*).

Approval duration: 6 months

B. Chronic Iron Overload due to Non-Transfusion Dependent Thalassemia Syndromes

(must meet all):

1. Diagnosis of chronic iron overload due to NTDT;
2. Age \geq 10 years;
3. Member must use generic deferasirox, unless contraindicated or clinically significant adverse effects are experienced;
4. Documentation of serum ferritin level $>$ 300 mcg/L and a LIC \geq 5 mg Fe/g dw;
5. Therapy does not include concurrent use of other iron chelators;
6. At the time of the request, member has none of the following contraindications:
 - a. GFR $<$ 40 mL/min/1.73 m²;
 - b. Platelet count $<$ 50 x 10⁹/L;
 - c. Severe hepatic impairment (Child-Pugh C);
7. Dose does not exceed the following (a or b):
 - a. Exjade: 20 mg/kg per day (*see Appendix D for dose rounding guidelines*);
 - b. Jadenu: 14 mg/kg per day (*see Appendix D for dose rounding guidelines*).

Approval duration: 6 months

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Chronic Iron Overload due to Blood Transfusions (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met all initial approval criteria;
2. Current documentation (within the past 30 days) shows a serum ferritin level \geq 500 mcg/L;
3. Therapy does not include concurrent use of other iron chelators;
4. If request is for a dose increase, new dose does not exceed the following (a or b):
 - a. Exjade: 40 mg per kg per day (*see Appendix D for dose rounding guidelines*);
 - b. Jadenu: 28 mg per kg per day (*see Appendix D for dose rounding guidelines*).

Approval duration: 12 months

B. Chronic Iron Overload due to Non Transfusion-Dependent Thalassemia Syndromes

(must meet all):

1. Currently receiving medication via Centene benefit or member has previously met all initial approval criteria;
2. Current documentation (serum ferritin within past 30 days; LIC within past 90 days) shows one of the following (a or b):

- a. If member has received < 6 months of Exjade/Jadenu, a serum ferritin level ≥ 300 mcg/L or an LIC ≥ 3 mg Fe/g dw;
- b. If member has received ≥ 6 months of Exjade/Jadenu, an LIC is ≥ 3 mg Fe/g dw;
3. Therapy does not include concurrent use of other iron chelators;
4. If request is for a dose increase, new dose does not exceed the following (a or b):
 - a. Exjade: 20 mg per kg per day (*see Appendix D for dose rounding guidelines*);
 - b. Jadenu: 14 mg per kg per day (*see Appendix D for dose rounding guidelines*).

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM. PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM. PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration
Fe/g dw: iron in milligrams per gram dry

GFR: glomerular filtration rate

LIC: liver iron concentration

NTDT: non-transfusion-dependent thalassemia

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Estimated GFR < 40 mL/min/1.73 m²
 - Poor performance status
 - High-risk myelodysplastic syndromes
 - Advanced malignancies
 - Platelet count < 50 x 10⁹/L
 - Known hypersensitivity to deferasirox or any component of Exjade of Jadenu
- Boxed warning(s):
 - Acute kidney injury, including acute renal failure requiring dialysis and renal tubular
 - Hepatic toxicity, including failure
 - Gastrointestinal hemorrhage

- Therapy requires close patient monitoring, including laboratory tests of renal and

*Appendix D: Dose Rounding Guidelines**

Weight-based Dose Range	Tablet for Oral Solution Quantity Recommendation
≤ 131.24 mg	125 mg tablet
131.25 mg – 262.49 mg	250 mg tablet
262.5 mg – 392.99 mg	125 mg tablet and 250 mg tablet
393 mg – 524.99 mg	500 mg tablet
525 mg – 655.99 mg	125 mg tablet and 500 mg tablet
656 mg – 787.49 mg	250 mg tablet and 500 mg tablet
787.5 mg – 917.99 mg	125 mg tablet, 250 mg tablet and 500 mg tablet
918 mg – 1,049.99 mg	2 x 500 mg tablets
1,050 mg – 1,180.99 mg	125 mg tablet and 2 x 500 mg tablets
1,181 mg – 1,312.49 mg	250 mg tablet and 2 x 500 mg tablets
1,312.5 mg – 1,442.99 mg	125 mg tablet, 250 mg tablet and 2 x 500 mg tablets
1,443 mg – 1,574.99 mg	3 x 500 mg tablets
Weight-based Dose Range	Oral Granules (sachets) Quantity Recommendation
≤ 94.49 mg	90 mg sachet
94.5 mg – 188.99 mg	180 mg sachet
189 mg – 283.49 mg	90 mg sachet and 180 mg sachet
283.5 mg – 377.99 mg	360 mg sachet
378 mg – 472.49 mg	90 mg sachet and 360 mg sachet
472.5 mg – 566.99 mg	180 mg sachet and 360 mg sachet
567 mg – 661.49 mg	90 mg sachet, 180 mg sachet and 360 mg sachet
661.5 mg – 755.99 mg	2 x 360 mg sachets
756 mg – 850.49 mg	90 mg sachet and 2 x 360 mg sachets
850.5 mg – 944.99 mg	180 mg sachet and 2 x 360 mg sachets
945 mg – 1,039.49 mg	90 mg sachet, 180 mg sachet and 2 x 360 mg sachets
1,039.5 mg – 1,133.99 mg	3 x 360 mg sachets

**This is part of a dose rounding guideline on select drug classes as part of an initiative conducted on a larger scale with multiple references and prescriber feedback.*

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Deferasirox (Exjade)	Transfusional iron overload	20 mg/kg body weight (calculate dose to the nearest whole tablet) PO QD	40 mg/kg/day
	NTDT syndromes	10 mg/kg body weight (calculate dose to the nearest whole tablet) PO QD	20 mg/kg/day
Deferasirox (Jadenu)	Transfusional iron overload	14 mg/kg body weight (calculated to nearest whole tablet/sachet) PO QD	28 mg/kg/day
	NTDT syndromes	7 mg/kg body weight (calculated to nearest whole tablet/sachet) PO QD	14 mg/kg/day

VI. Product Availability

Drug	Availability
Deferasirox (Exjade)	Tablets for oral suspension: 125 mg, 250 mg, 500 mg
Deferasirox (Jadenu)	Tablets/sprinkle (sachets): 90 mg, 180 mg, 360 mg

VII. References

1. Exjade Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2020. Available at <https://www.pharma.us.novartis.com/sites/www.pharma.us.novartis.com/files/exjade.pdf>. Accessed May 13, 2021.
2. Jadenu Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2020. Available at <https://www.pharma.us.novartis.com/sites/www.pharma.us.novartis.com/files/jadenu.pdf>. Accessed May 13, 2021.
3. Musallam KM, Angastiniotis M, Eleftheriou A, Porter JB. Cross-talk between available guidelines for the management of patients with beta-thalassemia major. *Acta Haematol.* 2013; 130: 64-73. DOI: 10.1159/000345734.
4. Hoffbrand AV, Taher A, Cappellini MD. How I treat transfusional iron overload. *Blood.* November 1, 2012; 120(18): 3657-3669.
5. Taher AT, Viprakasit V, Musallam KM, Cappellini MD. Treating iron overload in patients with non-transfusion-dependent thalassemia: Critical Review. *Am J Hematol.* 2013; 88: 409-415. DOI: 10.1002/ajh.23405.
6. Taher A, Musallam K, Cappellini MD. Guidelines for the management of non-transfusion dependent thalassaemia (NTDT) 2nd edition. Thalassaemia International Federation. 2018. TIF publication No. 22.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Converted to new template. Added weight-based max dose; safety criteria was applied according to the safety guidance discussed at CPAC and endorsed by Centene Medical Affairs. Updated references.	09.17	11.17
3Q 2018 annual review: policies combined for Centene Medicaid, HIM (new) and Commercial (new) lines of business; no significant changes; references reviewed and updated.	04.30.18	08.18
3Q 2019 annual review: contraindications caveat added to required Jadenu trial; the following contraindications are added: platelets, GFR; Child Pugh C restriction is removed; added requirement that member does not have severe hepatic impairment; references reviewed and updated. References reviewed and updated.	05.14.19	08.19
Added appendix D: dose rounding guidelines; added reference to appendix D within criteria.	03.05.20	05.20
3Q 2020 annual review: no significant changes; references reviewed and updated.	04.20.20	08.20

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Per July SDC and prior clinical guidance, modify redirection to require medical justification why generic deferasirox cannot be used for Exjade, brand Jadenu and Jadenu Sprinkle requests.	07.09.20	
3Q 2021 annual review: no significant changes; revised medical justification language for not using generic deferasirox to “must use” language; references for HIM line of business off-label use revised from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	05.16.21	08.21

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2015 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.