

## **Clinical Policy: Valbenazine (Ingrezza, Ingrezza Sprinkle)**

Reference Number: CP.CPA.364

Effective Date: 01.01.26

Last Review Date: 05.26

Line of Business: Commercial

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Valbenazine (Ingrezza<sup>®</sup>, Ingrezza<sup>®</sup> Sprinkle) is a vesicular monoamine transporter 2 (VMAT2) inhibitor.

### **FDA Approved Indication(s)**

Ingrezza and Ingrezza Sprinkle are indicated for the treatment of adults with

- Tardive dyskinesia (TD)
- Chorea associated with Huntington's disease

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results, or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Ingrezza and Ingrezza Sprinkle are **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Tardive Dyskinesia (must meet all):**

1. Diagnosis of TD secondary to a centrally acting dopamine receptor blocking agent (DRBA) (*see Appendix F*);
2. Prescribed by or in consultation with a psychiatrist or neurologist;
3. Age  $\geq$  18 years;
4. Ingrezza/Ingrezza Sprinkle is not prescribed concurrently with Austedo<sup>®</sup>/Austedo<sup>®</sup> XR or tetrabenazine;
5. Dose does not exceed both (a and b):
  - a. 80 mg per day;
  - b. 1 capsule per day.

**Approval duration: 12 months or duration of request, whichever is less**

##### **B. Chorea Associated with Huntington Disease (must meet all):**

1. Diagnosis of chorea associated with Huntington disease;
2. Prescribed by or in consultation with a neurologist;
3. Age  $\geq$  18 years;
4. Targeted mutation analysis demonstrates a cytosine-adenine-guanine (CAG) trinucleotide expansion of  $\geq$  36 repeats in the huntingtin (HTT) gene;

5. Evidence of chorea is supported by a Unified Huntington Disease Rating Scale (UHDRS) score ranging from 1 to 4 on any one of chorea items 1 through 7 (*see Appendix H*);
6. Ingrezza/Ingrezza Sprinkle is not prescribed concurrently with Austedo/Austedo XR or tetrabenazine;
7. Dose does not exceed both (a and b):
  - a. 80 mg per day;
  - b. 1 capsule per day.

**Approval duration: 12 months or duration of request, whichever is less**

**C. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial.

**II. Continued Therapy**

**A. All Indications in Section I (must meet all):**

1. Currently Member meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member meets one of the following (a or b):
  - a. For TD: Member is responding positively to therapy as evidenced by a reduction since baseline in any one of Abnormal Involuntary Movement Scale (AIMS) items 1 through 9 (*see Appendix G*);
  - b. For Huntington disease: Member is responding positively as evidenced by a reduction since baseline in any one of the UHDRS chorea items 1 through 7 (*see Appendix H*);
3. Ingrezza/Ingrezza Sprinkle is not prescribed concurrently with Austedo/Austedo XR or tetrabenazine;

4. If request is for a dose increase, new dose does not exceed both (a and b):
  - a. 80 mg per day;
  - b. 1 capsule per day.

**Approval duration: 12 months or duration of request, whichever is less**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

AIMS: Abnormal Involuntary Movement Scale

APA: American Psychiatry Association

CAG: cytosine-adenine-guanine

DRBA: dopamine receptor blocking agent

DSM-5-TR: Diagnostic and Statistical

Manual of Mental Disorders, Fifth Edition, Text Revision

FDA: Food and Drug Administration

HTT: huntingtin

TD: tardive dyskinesia

UHDRS: Unified Huntington Disease Rating Scale

VMAT2: vesicular monoamine transporter

*Appendix B: Therapeutic Alternatives*

Not applicable

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): known hypersensitivity to valbenazine or any components of Ingrezza or Ingrezza Sprinkle
- Boxed warning(s): depression and suicidal ideation and behavior in patients with Huntington's disease

*Appendix D: General Information - Tardive Dyskinesia*

- Ingrezza/Ingrezza Sprinkle should not be used concurrently with other VMAT2 inhibitors such as tetrabenazine or deutetabenazine as this is considered duplicate therapy.
- Medication-induced movement disorders, including tardive dyskinesia, are organized in the DSM-5-TR as follows: medication-induced parkinsonism, neuroleptic malignant syndrome, medication-induced acute dystonia, medication-induced acute akathisia, tardive dyskinesia, tardive dystonia/tardive akathisia, medication-induced postural tremor, other medication-induced movement disorder, antidepressant discontinuation syndrome, and other adverse effects of medication.
- Tardive dyskinesia is a type of movement disorder that occurs secondary to therapy with an antipsychotic medication or other DRBA (*see Appendix E*). (DSM-5-TR)
- Typical therapeutic drug classes containing DRBAs include first- and second-generation antipsychotics, antiemetics, and tri-cyclic antidepressants (*see Appendix F*). (DSM-5-TR)
- Other therapeutic drug classes containing agents that have been variously associated with movement disorders are listed below: (Waln 2013, Meyer 2014, Lerner 2015)
  - Antiarrhythmics
  - Antibiotics
  - Anticholinergics
  - Antidepressants
  - Antiepileptics
  - Antihistamines
  - Antimanics
  - Bronchodilators
  - Calcium channel blockers
  - Central nervous system stimulants
  - Dopamine agonists
  - Dopamine depleting agents
  - Dopaminergics
  - Glucocorticoids
  - Immunosuppressants
  - Mood stabilizers
  - Muscle relaxants
  - Oral contraceptives

*Appendix E: Tardive Dyskinesia: DSM-5-TR Definition*

**Tardive Dyskinesia (ICD-10 G24.01)**

- The essential features of tardive dyskinesia are abnormal, involuntary movements of the tongue, jaw, trunk, or extremities that develop in association with the use of medications that block postsynaptic dopamine receptors, such as first- and second-generation antipsychotic medications and other medications such as metoclopramide for gastrointestinal disorders. The movements are present over a period of  $\geq 4$  weeks and may be choreiform (rapid, jerky, nonrepetitive), athetoid (slow, sinuous, continual), or semirhythmic (e.g., stereotypies) in nature.
- Signs or symptoms of tardive dyskinesia develop during exposure to the antipsychotic medication or other dopamine blocking agent, or within 4 weeks of withdrawal from an oral agent (or within 8 weeks of withdrawal from a long-acting injectable agent). There must be a history of the use of the offending agent for  $\geq 3$  months (or 1 month in individuals age  $\geq 60$  years). Dyskinesia that emerges during withdrawal from an antipsychotic medication or other DRBA may remit with continued withdrawal from the medication. If the dyskinesia persists for  $\geq 4$  weeks, a diagnosis of tardive dyskinesia may be warranted..

*Appendix F: Centrally Acting Dopamine Receptor Blocking Agents*

Pharmacologic Class	Therapeutic Class		
	First-generation (typical) antipsychotics	Antiemetic agents	Tri-cyclic antidepressants
Phenothiazine	Chlorpromazine Fluphenazine Perphenazine Thioridazine Thiothixene Trifluoperazine	Chlorpromazine Perphenazine Prochlorperazine Promethazine* Thiethylperazine	Amoxapine <sup>†</sup>
Butyrophenone	Haloperidol	Droperidol Haloperidol**	
Substituted benzamide		Metoclopramide Trimethobenzamide	
Dibenzazepine	Loxapine		
Diphenylbutylpiperidine	Pimozide		
Pharmacologic Class	Second-generation (atypical) antipsychotics		
Quinolone	Aripiprazole, brexpiprazole		
Dibenzazepine	Asenapine		
Piperazine	Cariprazine		
Dibenzodiazepine	Clozapine, quetiapine		
Benzisoxazole	Iloperidone		
Benzisothiazole	Lurasidone, ziprasidone		
Thienobenzodiazepine	Olanzapine		
Pyrimidinone	Paliperidone, risperidone		

*(DSM-5-TR, Meyer 2014, Smith 2010, Clinical Pharmacology, Lexicomp)*

\*First generation H1 antagonist

\*\*Off-label use

<sup>†</sup>A dibenzoxapine that shares properties with phenothiazines

*Appendix G: The Abnormal Involuntary Movement Scale (AIMS)*

- The AIMS is a clinician-rated 12-item assessment tool developed by the National Institute of Mental Health to evaluate severity of involuntary movements in multiple movement disorders including TD. The AIMS is commonly used in both research and clinical practice.
- AIMS items 1-10 are rated on a 5-point scale (0 - none; 1 - minimal; 2 - mild; 3 - moderate; 4 - severe). Items 1-7 assess dyskinesia severity by body region (items 1-4 orofacial; items 5-7 extremity and trunk). Items 8-10 assess overall severity, incapacitation, and patient awareness respectively - item 8 uses the highest score of any one of items 1-7. Items 11 (dental) and 12 (dentures) are yes/no questions which help characterize lip, jaw, and tongue movements.
- See Munetz 1988 for additional information about the AIMS.

*Appendix H: Chorea: The Unified Huntington Disease Rating Scale (UHDRS)*

- The UHDRS encompasses motor, behavioral, cognitive, and functional components for use in evaluating patients with Huntington disease and is commonly used in both research and clinical practice.
- The American Academy of Neurology (AAN) guidelines evaluating pharmacologic therapies for chorea associated with Huntington disease describe the chorea subscore of the UHDRS motor component as a rating of 7 body regions (facial, bucco-oral-lingual, trunk, extremities) on a five-point scale from 0 to 4 with 0 representing no chorea.
- See Huntington Study Group 1996 and Mestre et al. 2018 for additional information about the UHDRS.

(AAN Guidelines 2012, Huntington Study Group 1996, Mestre 2018)

**V. Dosage and Administration**

Drug Name	Indication	Dosing Regimen	Maximum Dose
Valbenazine (Ingrezza, Ingrezza Sprinkle)	TD	40 mg PO once daily; after a week, increase to the recommended dose of 80 mg one daily. A dosage of 40 mg or 60 mg once daily may be considered depending on response and tolerability.	80 mg/day
Valbenazine (Ingrezza, Ingrezza Sprinkle)	Chorea associated with Huntington's disease	40 mg PO once daily; increase the dose in 20 mg increments every two weeks to the recommended dose of 80 mg once daily. A dosage of 40 mg or 60 mg once daily may be considered depending on response and tolerability.	80 mg/day

**VI. Product Availability**

Drug Name	Availability
Valbenazine (Ingrezza)	Oral capsules: 40 mg, 60 mg, 80 mg
Valbenazine (Ingrezza Sprinkle)	Oral sprinkle capsules: 40 mg, 60 mg, 80 mg

**VII. References**

1. Ingrezza and Ingrezza Sprinkle Prescribing Information. San Diego, CA: Neurocrine Biosciences, Inc.; October 2025. Available at: <http://www.ingrezza.com>. Accessed February 23, 2026.
- Tardive Dyskinesia*
2. Factor S, Comella C, Correll C, et al. Efficacy of valbenazine (NBI-98854) in subjects with tardive dyskinesia: Results of a long-term study (KINECT 3 extension) (S56.005). *Neurology*. April 18, 2017; 88(16): S56.005.
  3. Hauser RA, Factor SA, Marder SR. KINECT 3: A phase 3 randomized, double-blind, placebo-controlled trial of valbenazine for tardive dyskinesia. *Am J Psychiatry*. May 1, 2017; 174(5): 476-484. doi: 10.1176/appi.ajp.2017.16091037. Epub 2017 Mar 21.
  4. Keepers GA, Fochtmann LJ, Anzia JM, et al. The American Psychiatric Association practice guideline for the treatment of patients with schizophrenia. 2020. Available at <https://www.psychiatry.org/psychiatrists/practice/clinical-practice-guidelines>.

5. Munetz MR, Sheldon B. How to examine patients using the abnormal involuntary movement scale. *Hospital and Community Psychiatry*. November 1988;39(11):1172-77.
6. Bhidayasiri R, Jitkriksadakul O, Friedman JH, Fahn S. Updating the recommendations for treatment of tardive syndromes: a systematic review of new evidence and practical treatment algorithm. *Journal of the Neurological Sciences*. 2018;389:67-75.
7. Bashir HH, Jankovic J. Treatment of Tardive Dyskinesia. *Neurol Clin*. 2020;38(2):379-396.
8. Waln O, Jankovic J. An update on tardive dyskinesia: from phenomenology to treatment. *Tremor Other Hyperkinet Mov (N Y)*. July 12, 2013;3:tre-03-161-4138-1. DOI:10.7916/D88P5Z71. Print 2013.
9. Witter DP, Holbert RC, Suryadevara U. Pharmacotherapy for the treatment of tardive dyskinesia in schizophrenia patients. *Expert Opin Pharmacother*. April 26, 2017. DOI:10.1080/14656566.2017.1323874. [Epub ahead of print.]
10. Meyer TA, Belson TE, McAllister R. Tardive dyskinesia: a distressing drug-induced movement disorder. *US Pharm*. 2014;39(1):HS13-HS16.
11. Lerner PP, Miodownik C, Lerner V. Tardive dyskinesia (syndrome): current concept and modern approaches to its management. *Psychiatry Clin Neurosci*. June 2015;69(6):321-34.
12. Smith HS, Cox LR, Smith BR. Dopamine receptor antagonists. *Annals of Palliative Medicine*. July 2012;1(2). DOI: 10.3978/j.issn.2224-5820.2012.07.09.
13. Medication-Induced Movement Disorders and Other Adverse Effects of Medication. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, Text Revision (DSM-5-TR)*. American Psychiatric Publishing. February 2025. Accessed February 24, 2025. doi:10.1176/appi.books.9780890425787.Medication\_Induced\_Movement\_Disorders.

#### Huntington Disease

14. Stimming EF, Claassen DO, et al. Safety and efficacy of valbenazine for the treatment of chorea associated with Huntington's disease (KINECT-HD): a phase 3, randomized, double-blind, placebo-controlled trial. *Lancet*. 2023;22(6):494-504.
15. Potter NT, Spector EB, Prior TW. Technical standards and guidelines for Huntington disease testing. *Genet Med*. 2004;6(1):61-65.
16. Bean L and Bayrak-Toydemir R. American College of Medical Genetics and Genomics Standards and Guidelines for Clinical Genetics Laboratories, 2014 edition: technical standards and guidelines for Huntington disease. *Genet Med*. 2014 Dec;16(12):e2.
17. Bean L, Bayrak-Toydemir P, ACMG Laboratory Quality Assurance Committee. Addendum: American College of Medical Genetics and Genomics Standards and Guidelines for Clinical Genetics Laboratories, 2014 edition: technical standards and guidelines for Huntington disease. *Genet Med*. 2021;23(12):2461.
18. Kremer B, Goldberg P, Andrew SE. A worldwide study of the Huntington's disease mutation: the sensitivity and specificity of measure CAG repeats. *NEJM*. May 19, 1994; 330(20):1401-1406.
19. Armstrong MJ, Miyasaki JM. Evidence-based guideline: pharmacologic treatment of chorea in Huntington disease: report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology*. August 7, 2012;79:598-603.
20. Unified Huntington's disease rating scale: reliability and consistency. *Movement Disorder Society. Movement Disorders*. 1996;11(2):136-143.

21. Mestre TA, Forjaz MJ, Mahlknecht P, et al. Rating scales for motor symptoms and signs in Huntington’s disease: Critique and recommendation. International Parkinson and Movement Disorders Society. Movement Disorders Clinical Practice. 2018;5(2):111-117.  
DOI:10.1002/mdc3.1257.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created per SDC request (adapted from CP.PCH.48 which is being retired as HIM line of business is being added to CP.PHAR.340).	09.09.25	12.26
2Q 2026 annual review: no significant changes; references reviewed and updated.	02.23.26	05.26

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2026 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.