

Clinical Policy: Capivasertib (Truqap)

Reference Number: CP.PHAR.663

Effective Date: 03.01.24

Last Review Date: 02.26

Line of Business: Commercial, HIM/ICHRA, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Capivasertib (Truqap™) is a kinase inhibitor.

FDA Approved Indication(s)

Truqap is indicated:

- In combination with fulvestrant for the treatment of adult patients with hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer with one or more *PIK3CA/AKT1/PTEN*-alterations as detected by an FDA-approved test following progression on at least one endocrine-based regimen in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy;
- In combination with abiraterone and prednisone for the treatment of adult patients with metastatic androgen pathway modulation-naïve or -sensitive (APMN/S) prostate cancer that is PTEN-deficient as detected by an FDA-authorized test.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Truqap is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Breast Cancer (must meet all):**

1. Diagnosis of breast cancer;
2. Prescribed by or in consultation with an oncologist;
3. Age $\geq$ 18 years;
4. Disease has all of the following characteristics (a, b, c, and d):
 - a. HR-positive (i.e., estrogen or progesterone receptor [ER/PR]-positive);
 - b. HER2-negative;
 - c. Recurrent unresectable (local or regional) or metastatic;
 - d. Positive for *PIK3CA/AKT1/PTEN*-alteration(s);
5. Truqap is prescribed in combination with fulvestrant;
6. Member meets one of the following (a or b):
 - a. Disease progression or recurrence after an endocrine-based regimen (*see Appendix B*);
 - b. Recurrence on or within 12 months of completing adjuvant therapy (*see Appendix B*);

7. If member is a premenopausal female, member has been treated with ovarian ablation or is receiving ovarian suppression (*see Appendix D*);
8. For Truqap requests, member must use generic capivasertib, if available, unless contraindicated or clinically significant adverse effects are experienced;
9. Request meets one of the following (a or b):*
 - a. Dose does not exceed both of the following for 4 days followed by 3 days off (i and ii):
 - i. 800 mg per day;
 - ii. 4 tablets per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Prostate Cancer (must meet all):

1. Diagnosis of metastatic prostate cancer;
2. Prescribed by or in consultation with an oncologist or urologist;
3. Age $\geq$ 18 years;
4. Disease has both of the following characteristics (a and b):
 - a. APMN/S (previously referred to as “hormone-sensitive”);
 - b. PTEN-deficient ($\geq$ 90% of viable malignant cells with no specific cytoplasmic staining);
5. Member will use a gonadotropin-releasing hormone (GnRH) analog concurrently or has had a bilateral orchiectomy (*see Appendix D*);
6. Truqap is prescribed in combination with abiraterone and prednisone;
7. For Truqap requests, member must use generic capivasertib, if available, unless contraindicated or clinically significant adverse effects are experienced;
8. Request meets one of the following (a or b):*
 - a. Dose does not exceed both of the following for 4 days followed by 3 days off (i and ii):
 - i. 800 mg per day;
 - ii. 4 tablets per day;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the

relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Truqap for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively;
3. For Truqap requests, member must use generic capivasertib, if available, unless contraindicated or clinically significant adverse effects are experienced;
4. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed both of the following for 4 days followed by 3 days off (i and ii):
 - i. 800 mg per day;
 - ii. 4 tablets per day;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace/ICHRA, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace/ICHRA) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace/ICHRA, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace/ICHRA, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

APMN/S: androgen pathway modulation-naïve or -sensitive	HR: hormone receptor
ER: estrogen receptor	LHRH: luteinizing hormone-releasing hormone
FDA: Food and Drug Administration	NCCN: National Comprehensive Cancer Network
HER2: human epidermal growth factor receptor 2	PR: progesterone receptor
GnRH: gonadotropin-releasing hormone	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Endocrine therapy		
anastrozole (Arimidex [®])	1 mg PO QD	1 mg/day
exemestane (Aromasin [®])	25 mg PO QD	25 mg/day
toremifene (Fareston [®])	60 mg PO QD	60 mg/day
letrozole (Femara [®])	2.5 mg PO QD	2.5 mg/day
tamoxifen (Nolvadex [®] , Soltamox [®])	20 to 40 mg PO QD	40 mg/day
megestrol acetate	40 mg PO QID	160 mg/day
NCCN-recommended adjuvant therapy for HR-positive, HER2-negative disease		
CDK4/6 inhibitors to be used in combination with an aromatase inhibitor: Kisqali [®] (ribociclib) Verzenio [®] (abemaciclib) Ibrance [®] (palbociclib)	Varies	Varies

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): severe hypersensitivity to Truqap or any of its components
- Boxed warning(s): none reported

Appendix D: General Information

- NCCN recommendations in breast cancer
 - Ovarian ablation may be achieved with surgical oophorectomy or ovarian irradiation. Ovarian suppression utilizes luteinizing hormone-releasing hormone (LHRH)

- agonists that result in suppression of luteinizing hormone and release of follicle-stimulating hormone from pituitary and reduction in ovarian estrogen production. LHRH agonists include goserelin and leuprolide.
- The NCCN recommends that men with breast cancer be treated similarly to postmenopausal women, except that the use of an aromatase inhibitor is ineffective without concomitant suppression of testicular steroidogenesis.
- Examples of androgen deprivation therapy for prostate cancer include:
 - Bilateral orchiectomy (surgical castration)
 - LHRH agonist given with or without an anti-androgen:
 - LHRH agonists: Zoladex[®] (goserelin), leuprolide (Lupron Depot[®] or Eligard[®]), Trelstar[®] (triptorelin)
 - Anti-androgens: bicalutamide (Casodex[®]), flutamide (Eulexin[®]), nilutamide (Nilandron[®]), Xtandi[®] (enzalutamide), Erleada[®] (apalutamide), Nubeqa[®] (darolutamide)
 - LHRH antagonists: Firmagon[®] (degarelix), Orgovyx[®] (relugolix)

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Breast cancer, prostate cancer	400 mg orally BID for 4 days followed by 3 days off	800 mg/day

VI. Product Availability

Oral tablets: 160 mg, 200 mg

VII. References

1. Truqap Prescribing Information. Wilmington, DE: AstraZeneca; June 2026. Available at: <https://www.truqap.com/>. Accessed June 22, 2026.
2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at http://www.nccn.org/professionals/drug_compendium. Accessed June 22, 2026.
3. National Comprehensive Cancer Network. Breast Cancer Version 5.2025. Available at: http://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed December 5, 2025.
4. National Comprehensive Cancer Network. Prostate Cancer Version 5.2026. Available at: http://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed June 22, 2026.
5. Turner NC, Oliveira M, Howell SJ, et al. Capiwasertib in hormone receptor-positive advanced breast cancer. N Engl J Med. 2023;388(22):2058-2070.
6. Fizazi K, Clarke NW, De Santis M, et al. Capiwasertib plus abiraterone in PTEN-deficient metastatic hormone-sensitive prostate cancer: CAPItello-281 phase III study. Ann Oncol. 2026 Jan;37(1):53-68. doi: 10.1016/j.annonc.2025.10.004.
7. Micromedex[®] Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed December 5, 2023.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	12.05.23	02.24

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2025 annual review: no significant changes; removed the Coding Implications section since this is an oral agent; references reviewed and updated.	12.02.24	02.25
1Q 2026 annual review: no significant changes; added use of Truqap for recurrence within 12 months of adjuvant therapy to align with its original FDA approval; updated initial auth duration from 6 months to 12 months; references reviewed and updated.	11.06.25	02.26
RT4: added criteria for the newly approved prostate cancer indication; added ICHRA line of business.	06.22.26	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2024 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.