

Clinical Policy: Sargramostim (Leukine)

Reference Number: CP.CPA.262

Effective Date: 11.16.16

Last Review Date: 08.19

Line of Business: Commercial

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Sargramostim (Leukine[®]) is a recombinant human granulocyte-macrophage colony stimulating factor (GM-CSF).

FDA Approved Indication(s)

Leukine is indicated:

- To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in adult patients 55 years and older with acute myeloid leukemia (AML);
- For the mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation in adult patients;
- For the acceleration of myeloid reconstitution following autologous peripheral blood progenitor cell (PBPC) or bone marrow transplantation in adult and pediatric patients 2 years of age and older with non-Hodgkin's lymphoma (NHL), acute lymphoblastic leukemia (ALL) and Hodgkin's lymphoma (HL);
- For the acceleration of myeloid reconstitution following allogeneic bone marrow transplantation in adult and pediatric patients 2 years of age and older;
- For treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic bone marrow transplantation in adult and pediatric patients 2 years of age and older;
- To increase survival in adult and pediatric patients from birth to 17 years of age acutely exposed to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Leukine is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Acute Myelogenous Leukemia (must meet all):

1. Diagnosis of AML;
2. Prescribed for use following induction therapy for AML;
3. Age \geq 55 years;
4. Failure of Neupogen[®] or Zarxio[®], unless contraindicated or clinically significant adverse effects are experienced;

**Prior authorization is required for Neupogen and Zarxio*

5. Dose does not exceed 250 mcg/m² IV daily.

Approval duration: 6 months or to the member's renewal date, whichever is longer

B. Peripheral Blood Progenitor Cell Collection and Transplantation (must meet all):

1. Prescribed for one of the following (a or b):
 - a. Mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation;
 - b. Following autologous PBPC transplantation in members with NHL, ALL, or HL for acceleration of myeloid reconstitution;
2. Age ≥ 2 years;
3. Failure of Neupogen or Zarxio, unless contraindicated or clinically significant adverse effects are experienced;

**Prior authorization is required for Neupogen and Zarxio*

4. Dose does not exceed 250 mcg/m² IV or SC daily.

Approval duration: 6 months or to the member's renewal date, whichever is longer

C. Bone Marrow Transplantation (must meet all):

1. Prescribed for use in one of the following settings (a, b, or c):
 - a. Following autologous BMT in members with NHL, ALL, or HL for acceleration of myeloid reconstitution;
 - b. Following allogeneic BMT for acceleration of myeloid reconstitution;
 - c. Following BMT where engraftment is delayed or has failed;
2. Age ≥ 2 years;
3. Failure of Neupogen or Zarxio, unless contraindicated or clinically significant adverse effects are experienced;

**Prior authorization is required for Neupogen and Zarxio*

4. Dose does not exceed 500 mcg/m² IV daily.

Approval duration: 6 months or to the member's renewal date, whichever is longer

D. Acute Radiation Syndrome (must meet all):

1. Prescribed for use following suspected or confirmed acute exposure to myelosuppressive doses of radiation;
 2. Failure of Neupogen or Zarxio, unless contraindicated or clinically significant adverse effects are experienced;
- *Prior authorization is required for Neupogen and Zarxio*
3. Dose does not exceed 12 mcg/kg SC daily (See Section V for specific weight-based dosing).

Approval duration: 6 months or to the member's renewal date, whichever is longer

E. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial.

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria.
2. Member is responding positively to therapy;
3. Dose does not exceed the FDA-approved maximum recommended dose for the relevant indication.

Approval duration: 6 months or to the member’s renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ALL: acute lymphoblastic leukemia	H-ARS: hematopoietic syndrome of acute radiation syndrome
AML: acute myelogenous leukemia	NHL: non-Hodgkin's lymphoma
BMT: bone marrow transplantation	PBPC: peripheral blood progenitor cell
FDA: Food and Drug Administration	
GM-CSF: granulocyte-macrophage colony stimulating factor	

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
Neupogen (filgrastim), Zarxio (filgrastim-sndz)	AML: 5 mcg/kg SC or IV QD BMT: 10 mcg/kg IV or SC infusion QD PBPC collection: 10 mcg/kg SC bolus or continuous infusion QD Acute Radiation Syndrome:	AML: 30 mcg/kg/day IV or 24 mcg/kg/day SC BMT, PBPC collection, Acute Radiation Syndrome: 10 mcg/kg/day

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
	10 mcg/kg SC QD	

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): history of serious allergic reactions, including anaphylaxis
- Boxed warning(s): none reported

Appendix D: General Information

- Because of potential sensitivity of rapidly dividing hematopoietic progenitor cells, Leukine should not be administered simultaneously with cytotoxic chemotherapy or radiotherapy or within 24 hours preceding or following chemotherapy or radiotherapy.
- Use Leukine with caution in patients with pre-existing fluid retention, pulmonary infiltrates, or congestive heart failure.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
AML	250 mcg/m ² /day IV over a 4 hour period approximately on day 11 or four days following the completion of induction chemotherapy	250 mcg/m ² IV daily
PBPC collection and transplantation	250 mcg/m ² /day administered IV over 24 hours or SC once daily	250 mcg/m ² IV or SC daily
Myeloid reconstitution after autologous or allogeneic BMT	250 mcg/m ² /day IV over a 2 hour period beginning two to four hours after bone marrow infusion, and not less than 24 hours after the last dose of chemotherapy or radiotherapy	500 mcg/m ² IV daily
BMT failure or engraftment delay	250 mcg/m ² /day for 14 days as a 2 hour IV infusion	500 mcg/m ² IV daily
Acute Radiation Syndrome	Weight-based dose SC QD: >40 kg: 7 mcg/kg 15 to 40 kg: 10 mcg/kg <15 kg: 12 mcg/kg	See dosing regimen

VI. Product Availability

Lyophilized powder: 250 mcg single-dose vial
Solution: 500 mcg/mL multiple-dose vial

VII. References

1. Leukine Prescribing Information. Bridgewater, NJ: Sanofi-Aventis U.S., LLC.; March 2018. Available at: www.leukine.com. Accessed May 15, 2019.

2. National Comprehensive Cancer Network. Myeloid Growth Factors Version 2.2019. Available at: http://www.nccn.org/professionals/physician_gls/pdf/myeloid_growth.pdf. Accessed May 2, 2018.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed May 15, 2019.
4. DRUGDEX® System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed May 15, 2019.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J2820	Injection, sargramostim (GM-CSF), 50 mcg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Converted to new template. Minor changes to verbiage and grammar. References updated.	06.16.17	11.17
3Q 2018 annual review: added new indication for acute radiation syndrome; added criteria sets for each FDA indication to align with other lines of business; references reviewed and updated.	05.02.18	08.18
3Q 2019 annual review: no significant changes; references reviewed and updated.	05.15.19	08.19

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of

insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.