

Clinical Policy: Amlodipine/Atorvastatin (Caduet)

Reference Number: CP.CPA.237

Effective Date: 11.16.16

Last Review Date: 11.17

Line of Business: Medicaid – Medi-Cal

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Amlodipine/atorvastatin (Caduet[®]) is a combination of amlodipine, a calcium channel blocker, and atorvastatin, a HMG CoA-reductase inhibitor.

FDA approved indication

Caduet is indicated in patients for whom treatment with both amlodipine and atorvastatin is appropriate.

Amlodipine is indicated for the treatment of:

- Treatment of hypertension, to lower blood pressure
- Coronary Artery Disease (CAD)
 - Symptomatic treatment of chronic stable angina
 - Treatment of confirmed or suspected vasospastic angina (Prinzmetal's or Variant Angina)
 - Angiographically documented CAD
 - To reduce the risk for hospitalization for angina and coronary revascularization procedure in patients with recently documented CAD by angiography and without heart failure or an ejection fraction <40%,

Atorvastatin is indicated for:

- Prevention of Cardiovascular Disease:
 - Reduce the risk of myocardial infarction, reduce the risk of stroke, and reduce the risk for revascularization procedures and angina in adult patients without clinically evident coronary heart disease, but with multiple risk factors for coronary heart disease such as age, smoking, hypertension, low high-density lipoprotein cholesterol (HDL-C), or a family history of early coronary heart disease
 - Reduce the risk of myocardial infarction and reduce the risk of stroke in patients with type 2 diabetes, and without clinically evident coronary heart disease, but with multiple risk factors for coronary heart disease such as retinopathy, albuminuria, smoking, or hypertension.
 - Reduce the risk of non-fatal myocardial infarction, reduce the risk of fatal and non-fatal stroke, reduce the risk for revascularization procedures, reduce the risk of hospitalization for congestive heart failure (CHF), and reduce the risk of angina in patients with clinically evident coronary heart disease
- Hyperlipidemia
 - Heterozygous Familial and Nonfamilial Hypercholesterolemia:

- As an adjunct to diet to reduce elevated total-cholesterol (C), low-density lipoprotein cholesterol (LDL-C), apolipoprotein B (apo B), and triglyceride (TG) levels and to increase HDL-C in patients with primary hypercholesterolemia (heterozygous familial and nonfamilial) and mixed dyslipidemia (Fredrickson types IIa and IIb)
- Elevated Serum TG (triglyceride) levels:
 - As an adjunct to diet for the treatment of patients with elevated serum TG levels (Fredrickson type IV)
- Primary Dysbetalipoproteinemia:
 - Treatment of patients with primary dysbetalipoproteinemia (Fredrickson type III) who do not respond adequately to diet
- Homozygous Familial Hypercholesterolemia:
 - Reduce total-C and LDL-C in patients with homozygous familial hypercholesterolemia as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) or if such treatments are unavailable
- Pediatric Patients:
 - As an adjunct to diet to reduce total-C, LDL-C, and apo B levels in boys and postmenarchal girls, 10 to 17 years of age, with heterozygous familial hypercholesterolemia if after an adequate trial of diet therapy the following findings are present:
 - LDL-C remains ≥ 190 mg/dL or
 - LDL-C remains ≥ 160 mg/dL and:
 - There is a positive family history of premature cardiovascular disease or
 - Two or more other cardiovascular disease (CVD) risk factors are present in the pediatric patients

Limitation of use: Atorvastatin has not been studied in conditions where the major lipoprotein abnormality is elevation of chylomicrons (Fredrickson Types I and V).

Policy/Criteria

Provider must submit documentation (which may include office chart notes and lab results) supporting that member has met all approval criteria

It is the policy of health plans affiliated with Centene Corporation[®] that Caduet is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. All FDA-Approved Indications (must meet all):

1. Diagnosis of one of the following:
 - a. Hypertension;
 - b. Chronic stable angina;
 - c. Confirmed or suspected vasospastic angina (Prinzmetal's or Variant Angina);
 - d. Coronary artery disease documented by angiography and without heart failure or an ejection fraction $< 40\%$
2. Diagnosis of hyperlipidemia or one of the diagnoses for which atorvastatin is FDA approved;

3. Medical justification by physician for not using separate agents: atorvastatin and amlodipine;
4. Failure to achieve National Cholesterol Education Program (NCEP) goals, unless contraindicated or clinically significant adverse effects are experienced, to one generic formulary statin (lovastatin, pravastatin, simvastatin, atorvastatin) followed by Vytorin® or Crestor®.
5. Dose does not exceed 80 mg/day of atorvastatin.

Approval duration: Length of Benefit

B. Other diagnoses/indications

1. Refer to CP.PMN.53 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized)

II. Continued Therapy

A. All FDA-Approved Indications (must meet all):

1. Currently receiving medication via a health plan affiliated with Centene Corporation or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed 80 mg/day of atorvastatin.

Approval duration: Length of Benefit

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via a health plan affiliated with Centene Corporation and documentation supports positive response to therapy.
Approval duration: Duration of request or 12 months (whichever is less); or
2. Refer to CP.PMN.53 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized)

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy – CP.PMN.53 or evidence of coverage documents

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

NCEP: National Cholesterol Education Program

LDL: Low-density Lipoprotein Cholesterol

CVD: Cardiovascular Disease

CHF: Congestive Heart Failure

CAD: Coronary Artery Disease

Appendix B: General Information - N/A

Appendix C: Therapeutic Alternatives

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
atorvastatin (Lipitor [®])	10 - 80 mg PO QD	80 mg daily
amlodipine (Norvasc [®])	2.5 - 10 mg PO QD	10 mg daily
lovastatin (Mevacor [®])	10-80 mg PO QD or BID	80 mg daily
pravastatin (Pravachol [®])	10 – 80 mg PO QD	80 mg daily
simvastatin (Zocor [®])*	5 - 40 mg PO QD (Coverage of the 80 mg strength requires PA)	80 mg daily
ezetimibe/simvastatin (Vytorin [®] *)	10/10 mg - 10/80 mg PO QD (Coverage of the 10/80 mg strength requires PA)	10/80 mg daily
Rosuvastatin (Crestor [®] *)	5 - 40 mg/day PO QD	40 mg daily

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

* Requires Prior Authorization

V. Dosage and Administration

Drug Name	Dosing Regimen	Maximum Dose
Amlodipine/atorvastatin (Caduet)	2.5/10 mg PO QD Pediatric Dosing (age 10 - 17 years): 2.5/10 mg PO QD	Adult: 10/80mg PO QD Pediatric: 5/20mg PO QD

VI. Product Availability

Tablet: 2.5/10, 2.5/20, 2.5/40, 5/10, 5 /20, 5 /40, 5/80, 10/10, 10 /20, 10/40, 10/80 mg

VII. References

1. Caduet [Prescribing Information] New York, NY:Pfizer;March 2015.
2. Lipitor [Prescribing Information] New York, NY: Pfizer; March 2015.
3. Norvasc [Prescribing Information] New York, NY: Pfizer; March 2015.
4. Micromedex[®] Healthcare Series [Internet database]. Greenwood Village, CO: Thompson Healthcare. Updated periodically. Accessed January 14, 2016
5. Caduet. American Hospital Formulary Service Drug Information. Available at: <http://www.medicinescomplete.com/mc/ahfs/current/>. Accessed January 14, 2016.
6. Clinical Pharmacology Web Site. Available at <http://clinicalpharmacology-ip.com/>. Accessed January 14, 2016.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Converted to new template. Minor changes to verbiage and grammar. References updated.	06.14.17	11.17

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

CLINICAL POLICY
Amlodipine/Atorvastatin



Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.